**INDEX.**


A.

*Abramis brama*, 177
*Acanthocyclops*, 16
*robustus*, 70, 153, 163, 185
*Acanthodiaptomus denticornis*, 2, 7
*Aedes*
*aegypti*, 70
*vezans*, 54
African Lake district, 17
Aggregation, 163
Akinesis, 155
*Alburnus alburnus*, 42
Algeria, 63
Allometry, 41
*Alona*
*cf setulosa*, 73
*cf. cambouei*, 27
*diaphana*, 13, 24
Amazon river, 233
Ambush predator(s), 207, 223
*Ambystoma tigrinum*, 100
*Anisops*, 20
*Anopheles*
*freeborni*, 67
*stevensi*, 70
Anostraca, 68, 119
Anoxic water, 210
Anti-fish defense, 49
Anti-predator defense, 41
('neckteeth'), 51
*Arctodiaptomus spinosus*, 108
Argentina, 227
*Argyrodiaptomus furcatus*, 93
*Ascomorpha*, 128
*ecaudis*, 221
*Asplanchna*, 14, 16, 188, 127 163
*brightwelli*, 19
*girodi*, 18
*priodonta*, 2, 3, 19
*sieboldi*, 19
sp., 19, 205
Attack, 153
Australia, 28

B.

Balanced Predation Hypothesis, 215
Barra Bonita reservoir, 91, 141
*Bdellocephala punctata*, 62
Belgium, 154, 164
*Bosmina*, 27, 128, 163
*angularis*, 36
*coregoni*, 169, 212, 207, 227
*fatalis*, 34, 36
*hagmani*, 100
*longirostris*, 2, 34, 121, 133, 134, 186, 212
*Bosminopsis*, 27
*Brachionus*
*angularis*, 120
*calyciflorus*, 36, 141
spp., 34
*Branchinecta*
*coloradensis*, 67
*gigas*, 119
*paludosa*, 68, 70
*Branchipus schaefferi*, 69
Brazil, 141
*Bothromesostoma personatum*, 62
*Bythotrephes*, 14, 16, 205
*longimanus*, 107

C.

*Caenorhabditis elegans*, 121
Cannibalism, 230
in *Chaoborus*, 39
Capture probabilities, 212
Carapace strength/integrity, 153
Cascading effects of vertebrate and invertebrate predation, 9

Cellulose acetate
*Ceriodaphnia*, 27, 128
*cornuta*, 34, 141
*silvestris*, 93
*quadranglua*, 2
*Chaetogaster diaphanus*, 70
*Chaoborus*, 6, 13, 15, 22, 33, 54, 94, 107, 215, 234
*americanus*, 51, 180
*cristallinus*, 6
*flavicans*, 33, 67, 206, 227
*obscuripes*, 206
*punctipennis*, 215
sp., 70, 205
*trivittatus*, 67
Cheat Lake, 133
Chemical defenses, 228
*Chilodonella sp.*, 120
Chironomid larvae, 61
*Chironomus riparius*, 68
*Chydorus sphaericus*, 2, 65, 103, 163
Clearance rate, 121
Coexistence, 185
Coleoptera, 23
Community structure, 61
*Conochilus unicornis*, 2
Copepod demographic structure, 165
Copepodite development times, 189
*Coregonus albula*, 42
*Cryptomonas*, 224
*Culex tarsalis*, 67, 92
*Culiseta melanura*, 70
Cyclomorphosis in *Daphnia*, 41
Cyclopoid predation, 153, 163
*Cyclops*, 16
*strenuus*, 121, 227
*vicinus*, 34, 187
*vicinus vicinus*, 2, 4, 70

D.

*Daphnia*, 27, 103, 129, 154
*ambigua*, 34, 138, 166
*coloradensis*, 67
*cristata*, 43
*cucullata*, 41, 186
*curvirostris*, 27
*galeata*, 187
*hyalina*, 41, 42, 166
*laevis*, 92, 227
*longispina*, 2
*magna*, 63, 79, 121
*middendorffiana*, 67, 91
*parvula*, 133, 139
*pulex*, 51, 54, 67, 92
*similis*, 63
Dayat Ifrah, 92
Dead-man response, 155
Demographic data, 1
*Dendrocoelum lacteum*, 62
Density risk (DR), 205
Developmental bottlenecks, 205
*Diacyclops*, 16
Diapause, 5, 189
*Diaphanosoma*, 27
*brachyurum*, 34, 100, 107, 108, 138, 187
*leuchtenbergianum*, 139
*Diaptomus arcticus*, 7, 68
*floridanus*, 138
*kenai*, 207
*pallidus*, 133, 138
*shoshone*, 7, 67
*Dicranophorus*, 16
Direct observations, 1
Diurnal migration, 163
Diversity, 14
of tropical zooplankton, 13
*Dugesia dorotocephala*, 70
*Dunhevedia crassa*, 73
Dystiscidae, 16, 23

E.

Effect of temperature on handling time, 155
Effect of water turbulence on vertical distribution, 181
Egg vulnerability, 153
Electrophoretic phenotypes, 52
Embit, 164
Embryonic development times, 200

Enclosures, 215
Encounter probability, 207
*Eodiaptomus japanicus*, 34
*Epischura nevadensis*, 138
Escape behaviour, 99
response, 153
tactics, 215, 224
Ethiopian rift valley lakes, 17
*Eucyclops*, 16
*Eudiaptomus*, 58
*gracilis*, 207
Exclosure treatment, 21

F.

FRG, 185
Facultative predators, 15
Fecundity of *C. vicinus vicinus*, 5
Feeding preferences of *Mesostoma*, 94
Field experiments, 235
*Filinia*
*hofmanni*, 2
*longiseta*, 120
Filter feeding, 119, 210
Fluorescamine, 56
Food preference, 194
France, 1, 2

G.

*Gonyostomum semen*, 217
Grazing, 2
Guelta Adessei, 63
Gut analysis of late copepodite stages, 5
contents, 1

H.

Handling time, 153
Helmets, 41
Hemiptera, 20
*Hesperodiaptomus*, 16
*Heterocope*, 16
*Hexarthra mira*, 36, 38, 111
Horizontal distribution, 163
*Hydra*, 15, 26
Hydracarina, 13, 16, 230
Hydrophilidae, 16

I.

Index of Dispersion, 166
Indonesia, 63
Ingestion rates, 185, 194
Invertebrate predators, 9
Isometric growth, 41

J.

Japan, 34
Juvenile predators, 153

K.

Kairomone, 51
*Kellicottia*, 129
*bostoniensis*, 215, 221
*Keratella*
*cochlearis*, 2, 154, 163, 166
*longispina*, 2
*quadrata*, 2, 120, 121
spp., 34
*taurocephala*, 223
*valga*, 36, 38

L.

Laguna Aeroclub, 227
Lake Awasa, 19
Aydat, 1, 2
Dom Helvecio, 92
Donk, 154, 164
Gardsjon, 205
Glubokoe, 227
Jacare, 65
Kariba, 22
Kasumigaura, 34
Kivu, 17, 27
Lanao, 22

Majcz, 41
Mikolajskie, 41
Ontario, 22
Ranao, 63
Toba, 22
Turkana, 22
Valencia, 22
Zwai, 19
*Lepomis*
*cyanellus*, 71
*macrochirus*, 127
*Leptestheriella*, 119
*Leptodora*, 14, 16
*kindti*, 42, 107, 227
Lime treatment, 207
Liming, 205
Limitation 'from the top', 207
Littoral, 13, 61
Lloyd's measure of patchiness, 166
*Loxodes* sp., 120

M.

*Macrocyclops*, 16
Malaysia, 18
*Mallomonas sp.*, 121
Malpighian tubules, 53
Mechanoreceptors, 107
*Megacyclops*, 16
Mesocosms, 235
Mesocosm experiment, 217
*Mesocyclops*, 16, 19, 21
*aspericornis*, 19
*brasilianus*, 142
*edax*, 125, 127, 133
*gigas*, 147
*kieferi*, 141
*leuckarti*, 70, 185
*longisetus*, 70, 94, 141
*mendocinus*, 142
*pehpeiensis*, 19
*splendidus*, 19
*Mesostoma*, 103
*appinum*, 70
*ehrenbergi*, 67, 91
*lingua*, 63, 79, 92
*macroprostatum*, 71
sp., 61, 91
*timbuke*, 70
*zariae*, 70, 71
*Metacyclops*, 142
*Microcyclops*, 16
*Micropterus salmoides*, 127
Middle Tiete river, 141
*Moenkhausia intermedia*, 227
*Moina*, 103
*brachiata*, 188
*micrura*, 34, 36, 73
sp., 65
Morocco, 63
Morphogen, 51, 52
Morphological variation, 41
Mortality rates of
*K. cochlearis*, 4
Mosquito larvae, 61
Mosquitoes, 61
Mosquitofish (*Gambusia affinis*), 71
Mucus trapping, 61
*Mysis*, 16
*relicta*, 107, 125, 180

N.

*Naucoris scutellaris*, 19, 20
*Neomysis*, 16, 33
*intermedia*, 34
Neurotoxins, 75
Neusiedler See, 107
North temperate zone, 13
*Notodiaptomus iheringi*, 149
Notonectidae, 23
Numerical response, 227, 235

O.

Obligate predators, 15
Odonata larvae, 16, 26
Ontario, 18
Optimum foraging, 75
*Oscillatoria*
*redeckei*, 186
sp., 121
*tenuicaudis*, 73
*Osmerus eperlanus* (L.), 42
Oxygen depletion, 200

P.

*Parabroteus*, 16
Paralyzing toxin, 67
Patchiness, 164
*Pediastrum sp.*, 121
Pelagic area, 2
    flatworms, 63
    zone, 33
*Pelecus cultratus*, 115
Pennsylvania, 127
*Perca fluvialitis*, 177
*Phaenacera* species, 69
*Physiocypria*, 93
Piona predation rates, 227, 230
Plankton community structure, 133
*Plea*, 20
Pleiidae, 23
*Ploesoma*, 14, 16
Poland, 41
*Polyarthra*
    *dolichoptera*, 2
    *remata*, 2, 128
    sp., 34, 187
    *trigla*, 38
    *vulgaris*, 163, 215, 221
*Polyphemus*, 14, 16, 205
    *pediculus*, 107
*Pompholyx*
    *complanata*, 38
    *sulcata*, 163
Population dynamics, 2
Potential predation pressure, 235
Predation
    cycle, 153
    effects, 33
    effects by *Chaoborus*, 35
    on soft-bodied rotifers, 223
    rate, 107, 143
    risk, 125
    sequence, 8
Predatory Cladocera, 13
    behavior, 234
    losses, 215
Prey community structure, 205
    handling, 234
    pursuit, 234
    recognition, 234
    selection, 107, 133
    selection tactics, 207
    spectrum of *M. leuckarti*, 200
    vulnerability, 125
Professoren-Weiher, 185
*Pseudodiaptomus inopinus*, 34

R.

Refuges, 235
Resting stages, 191, 234
Rhabdocoeles, 26
*Rhynchomesostoma rostratum*, 92
Rice Fields, 19
Rice field enclosures, 71
Rio Doce Valley, 63, 91
Ruanda, 63

S.

*Scardinius erythrophtalamus*, 187
*Scenedesmus* sp., 121
Seasonal changes, 34
    in the abundance of *Neomysis intermedia*, 35
Seasonal succession, 1
Seasonal variations in density, 3
Sex ratio, 191
Sex specific mortality, 185
Sexual size dimorphism, 198
*Simocephalus*, 26
    *vetulus*, 92
Sit-and-wait predation, 61
Size variation, 191
Southern hemisphere, 234
Spatial overlap, 163, 235
Spatio-temporal distribution analysis, 1
Sri Lanka, 18
Stability, 234
*Stenostonum sthenum*, 70
*Stizostedion lucioperca*, 177
*Streptocephalus*
    *dichotomus*, 119
    *proboscideus*, 119

*Streptomyces griseus* Pronase, 53
Successive presence, 9
Summer decline of rotifers, 9
Summer depression in zooplankton, 227
Sweden, 205
Swimming velocities, 207
*Synchaeta*
*kitina*, 154
*oblonga*, 215, 221
*pectinata*, 154

T.

Takahamairi Bay, 34
Tanganyika, 17
Temperate zone, 1, 28, 233
Tender Bog Lake, 52
*Tetrahymena pyriformis*, 121
*Thermocyclops*, 15, 21, 142
*crassus*, 187
*decipiens*, 142
*minutus*, 93, 142
*taihokuensis*, 34
Top-down control, 215
Triangle Lake, 218
*Trichocerca cylindrica*, 2
Triclad flatworms, 62
*Triturus*, 70
Tropical lakes, 233
region, 14
zooplankton, 13
Tropics, 28
*Tropocyclops*, 129
Typhloplanid flatworms, 61

V.

Vertebrate predation, 2
Vertical distribution, 103, 163
of zooplankton, 96
Vertical migration, 91, 125
of *Mesostoma* sp., 98
Vertical overlap, 130
Vertical segregation between predator and prey, 163
*Volvox*, 119
Vulnerability, 44, 215
to predation, 41, 43

W.

Water mites, 227
Whole-community biomanipulations, 224

Z.

Zooplankton, 14
community maturity, 28
community structure, 33, 125, 215